




Key Fill Vaporiser Filling & Draining Instructions

Keyfiller - Filling

- Warning: Fill the vaporiser only when the control dial is in the OFF position and do not turn the dial on during filling or attempt to fill beyond the 'FULL' mark.
- Caution: Always hold the bottle below the level of the end of the filling adaptor and also below the level of the filler port until the adaptor is clamped in position.
- Caution: Turn screw slowly if vaporiser is pressurised. Turn top retaining screw on Vaporiser Filler unit counterclockwise and take out Filler Plug.



- 1. Insert the end of the Adaptor with the two holes downwards into the top filler port, take care to bend Tube down slightly so bottle is below inlet level.
- 2. After insertion, turn the retaining screw clockwise and tighten to seal Filler Adaptor in the Filler Socket.
- 3. Raise the bottle above the level of the filler port now and begin filling.

 Note: The stream of bubbles inside the bottle will confirm that filling is taking place. The Vaporiser will fill in 20-30 secs.

 When the vaporiser is filled to the full level lower the Bottle and remove the Adaptor.
- 4. Replace the Filler Plug and tighten the Top retaining screw again.

The Vaporiser is now ready for use

Liquid Capacity Amount of Anaesthetic Agent to fully charge the vaporiser = 150ml (nominal) Amount retained by Wick System = 35ml (nominal)

During initial fill, level may not begin to show until the wick is fully saturated









Draining



- 1. Unscrew the bottom retaining screw and insert the Adaptor into the bottom socket and tighten again.
- 2. Ensure that the bottle is held below the level of the end of the Adaptor and open the Drain Valve.

 Note: To let air vent from the bottle remove the filler plug.
- 3. When the vaporiser is drained remove Adaptor and tighten the Drain Valve and bottom retaining screw.





